

---

# MYCOTAXON

Volume 116, pp. 183–189

DOI: 10.5248/116.183

April–June 2011

---

## Hydnoid basidiomycetes new to Brazil

ALICE DA CRUZ LIMA GERLACH & CLARICE LOGUERCIO-LEITE

*Departamento de Botânica, Centro de Ciências Biológicas, Universidade Federal de Santa Catarina, Campus Universitário, 88040-900, Florianópolis, SC, Brazil*

CORRESPONDENCE TO: [alice\\_gerlach@yahoo.com.br](mailto:alice_gerlach@yahoo.com.br) & [clleite@ccb.ufsc.br](mailto:clleite@ccb.ufsc.br)

**ABSTRACT** — A survey of wood-decaying fungi from an *Araucaria* forest in the state of Santa Catarina in southern Brazil yielded numerous species of *Agaricomycetes*. Three hydneous species collected (*Mycobonia brunneoleuca*, *Mycoacia aurea* and *Spongipellis africana*) represent first records from Brazil. Illustrations and keys to the Brazilian species of *Mycobonia*, *Mycoacia*, and *Spongipellis* are provided.

**KEY WORDS** — *Polyporales*, corticioid, fungal distribution

### Introduction

The Atlantic Forest is still common in the state of Santa Catarina, where secondary forest dominates much of the landscape and 23% of the original vegetation cover remains (SOS Mata Atlântica/INPE 2009). The most outstanding feature of this subtropical region is the large extent of mixed *Araucaria* forests that cover the inner plateaus of southern Brazil and the Misiones Province in Argentina (Oliveira-Filho et al. 2009). These forests are easily recognized by their canopies, which are dominated by the chandelier-like crowns of *Araucaria angustifolia* (Bertol.) Kuntze (Sonogo et al. 2007), and form complex mosaics with grasslands at higher altitudes (Jarenkow & Budke 2009).

Previous surveys and reviews of hydneous species from Brazil were published by Rick (1932a,b, 1959), Bononi (1979, 1981), Bononi et al. (1981, 2008), Hjortstam & Bononi (1986a,b, 1987), Capelari & Maziero (1988), Sótão et al. (1991), Jesus (1993), Nietiedt & Guerrero (2000), Gibertoni et al. (2007), and Baltazar & Gibertoni (2009).

The aim of this study was to increase knowledge of hydneous basidiomycetes (*Agaricomycetes*) from *Araucaria* forests in Brazil.

## Materials & methods

The study was made in the Reserva Particular do Patrimônio Natural Rio das Furnas (RPPN Rio das Furnas) located in the Alfredo Wagner municipality, in Santa Catarina, Brazil (27°40'45"S, 49°10'38.4"W). This is a 10 ha reserve within the *Araucaria* forest region.

Microscopic characters were examined and measured using light microscopy, on slides mounted with 1% aqueous phloxine solution (plus 1% or 5% KOH) and Melzer's reagent (Ryvarden 1991). Drawings were made with the aid of a drawing tube. Voucher specimens are archived at FLOR (Thiers 2007).

## Taxonomy

*Mycobonia brunneoleuca*, *Mycoacia aurea*, and *Spongipellis africana* represent new records from Brazil and are described below based on the newly collected material. Keys including related Brazilian taxa are also provided.

### *Gloeophyllaceae* Jülich

*Mycobonia brunneoleuca* (Berk. & M.A. Curtis) Pat., *Essai Tax. Hyménomyc.*: 75. 1900. FIG. 1

VOUCHER MATERIAL: BRAZIL. SANTA CATARINA: Alfredo Wagner, RPPN Rio das Furnas, 07.VII.2008, Gerlach & Giovanka 122, FLOR 32323.

BRIEF DIAGNOSIS: *Mycobonia brunneoleuca* is characterized by the longer hyphal pegs ( $\leq 180 \mu\text{m}$  outside the hymenium), robust basidia ( $\leq 80 \times 13\text{--}20 \mu\text{m}$ ), and ovoid-elliptic basidiospores,  $15\text{--}24(-25) \times (6\text{--})7\text{--}11 \mu\text{m}$ , which contain many oil droplets.

ADDITIONAL SPECIMENS EXAMINED — BRAZIL. SANTA CATARINA: Itapiranga, Scholz s/n, 15.IV.1985, FLOR 0109; Três Barras, Flona, Drechsler-Santos s/n, 22.XI.2003, FLOR 31465; Mondáí, Linha Uruguai, 27.XII.2006, Campos-Santana, Santana & Souza-Rodrigues 188, FLOR 32226; 23.V.2007 Campos-Santana & Santana 223, FLOR 32227.

DISTRIBUTION: Neotropical.

COMMENTS: *MYCOBONIA BRUNNEOLEUCA* AND *M. FLAVA* (Sw.) Pat. are macroscopically indistinguishable except for their basidiospores, which are wider and ovoid-elliptic in *M. brunneoleuca*. Dennis (1970), who regarded these taxa as the same species, considered them common in cloud forests of Venezuela (e.g., Sierra de la Costa) between 1600 and 2000 meters. According to Reid (1976) it is possible that both species occupy the same geographical region but live in different niches, with *M. brunneoleuca* occurring at higher elevations. *Mycobonia flava* was previously collected within the Mondáí (220 m elev.) and Itapiranga ( $\leq 544$  m) municipalities and the Floresta Nacional de Três Barras reserve (766 m) in Santa Catarina. Our material was collected between 750 and 900 m in *Araucaria* forest, which agrees with observations made by Reid.




FIG. 1. *Mycobonia brunneoleuca*: A, hyphal pegs; B, basidiospores; C, basidia.  
Scale bars = 10  $\mu\text{m}$ .

#### Key to *Mycobonia* species in Brazil

1. Spores ovoid-ellipsoid, (6–)7–10  $\mu\text{m}$  wide ..... *M. brunneoleuca*  
 1. Spores fusoid-ellipsoid, 5–7  $\mu\text{m}$  wide ..... *M. flava*

#### *Meruliaceae* P. Karst.

*Mycoacia aurea* (Fr.) J. Erikss. & Ryvarde, Corticiac. N. Europe 4: 877. 1976. FIG. 2

VOUCHER MATERIAL: BRAZIL. SANTA CATARINA: Alfredo Wagner, RPPN Rio das Furnas, Reck, 17.VII.2009, Gutjahr 09, FLOR 32415.

BRIEF DIAGNOSIS: *Mycoacia aurea* has resupinate basidiomata that are very soft when fresh and with a horny consistency when dried. The species is characterized by a palisade hymenium ( $\leq 35 \mu\text{m}$  thick), subiculum with thin to slightly thick-walled, clamped generative hyphae, clavate 4-spored basidia with basal clamps, and suballantoid basidiospores ( $4\text{--}5 \times 1.5\text{--}2 \mu\text{m}$ ).

ADDITIONAL SPECIMEN EXAMINED: SWEDEN. DARLNA: Husby, Rankholmen i Flinesjön, naturreservat, 6,5 Km SO Husby K:a. Lat/long: 60°22'N, 16°06' E, Grid:




FIG. 2. *Mycoacia aurea*: A, hymenium; B, basidiospores; C, generative hyphae.  
Scale bars = 10 µm.

RN 6694300 1517200, Alt. 80 m. Prunus-Laga i graalskog, alvplan, 18.III.1990, Janolof Hermansson S253, Herb. Univ. Upsaliensis F-140874, 425667.

DISTRIBUTION: Cosmopolitan.

COMMENTS: Eriksson & Ryvar den (1976) distinguish *M. aurea* from other *Mycoacia* species by its small allantoid basidiospores, absence of cystidia, and absence of a red KOH reaction. Hjortstam & Ryvar den (1996) noted that *M. aurea* lacks incrust ed hyphal endings in its spines and has suballantoid basidiospores, which coincides with our material. The specimen from Sweden (F-140874) has some encrust ed hyphae but is otherwise very similar to the Brazilian specimen.

**Key to *Mycoacia* species in Brazil**

(based on Hjortstam & Ryvar den 1996)

- 1. Hymenophore tuberculate, subulate cystidia rare ..... *M. livida*
- 1. Hymenophore hydroid, cystidia either not subulate or absent ..... 2

2. Cystidia present .....3  
 2. Cystidia absent .....4  
 3. Cystidia subulate to acicular, encrusted hyphae present ..... *M. fuscoatra*  
 3. Cystidia subfusiform, encrusted hyphae absent ..... *M. uda*  
 4. Basidiospores allantoid,  $4-4.5 \times 1-1.25 \mu\text{m}$  ..... *M. subconspersa*  
 4. Basidiospores suballantoid,  $3.5-5.5 \times 1.5-2 \mu\text{m}$  ..... *M. aurea*

### *Polyporaceae* Corda

*Spongipellis africana* Ipulet & Ryvarden, Syn. Fung. 20: 97. 2005. FIG. 3

VOUCHER MATERIAL: BRAZIL. SANTA CATARINA: Alfredo Wagner, RPPN Rio das Furnas, 17.XII.2008, Gerlach, Giovanka 155, FLOR 32416.

BRIEF DIAGNOSIS: *Spongipellis africana* has widely adhered to effuse reflexed basidiomata that are soft when dried, clavate 4-spored basidia with basal clamps, and basidiospores that are (sub)globose [ $4-5(-6) \times 4-5 \mu\text{m}$ ], thick-walled, smooth, and with positive cyanophilic reaction.

ADDITIONAL SPECIMEN EXAMINED: URUGUAY. Dto Treinta e tres Quebrada de los Cuervos, on dicotyledonous dead wood, 1.VII.1993, C. Prigioni MVM 374, BAFC 34059.

DISTRIBUTION: Pantropical.




FIG. 3. *Spongipellis africana*: A, hymenium; B, basidiospores; C, cystidia.

Scale bars = 10  $\mu\text{m}$ .

COMMENTS: According to Ipulet & Ryvarden (2005), *S. africana* is macroscopically identical to *S. pachyodon* (Pers.) Kotl. & Pouzar but easily separated by its smaller basidiospores. Ipulet & Ryvarden (2005) did not observe

cystidia in this species; however, our material has inconspicuous metuloid cystidia in the hymenium. The specimen from Uruguay (BAFC 34059) was previously identified as *S. pachyodon* but it has smaller basidiospores. Therefore, we believe that this specimen is also *S. africana*, which extends the distribution of *S. africana* to Brazil and Uruguay.

#### Key to *Spongipellis* species in Brazil

(based on Piatek et al. 2004, Rajchenberg & Meijer 1990, Gilbertson & Ryvarden 1986)

1. Hymenophore distinctly hydroid. . . . . 2
1. Hymenophore poroid, or sometimes sinuous to daedaleoid . . . . . 3
2. Basidiospores globose to broadly ellipsoid,  $5.5\text{--}7 \times 4.5\text{--}5.5 \mu\text{m}$  . . . . . *S. pachyodon*
2. Basidiospores globose,  $4\text{--}4.5 \mu\text{m}$  . . . . . *S. africana*
3. Pores subgyrose, 1–3/mm, compressed when dry, dentate to fimbriate, hymenophore pinkish red to pinkish salmon when fresh, basidiospores ellipsoid to broadly ellipsoid,  $4.6\text{--}6 \times 3.3\text{--}4.3 \mu\text{m}$  . . . . . *S. caseosus*
3. Pores angular, 4–5/mm, hymenophore white to cream, basidiospores broadly ellipsoid to globose (ovoid),  $4.5\text{--}6 \times 4\text{--}5 \mu\text{m}$  . . . . . *S. fractipes*

#### Acknowledgments

The authors are grateful to Gabriela Giovanka, Renato Rizzaro, Matheus Reck and Márcio Gutjahr, especially for their help with the fieldwork. We are also grateful to the reviewers, Karen Nakasone (USA) and Maria Alice Neves (Brazil), as well Shaun Pennycook (New Zealand) and would like to thank them for their suggestions and for improving this article.

#### Literature cited

- Baltazar JM, Gibertoni TB. 2009. A checklist of the aphylloroid fungi (*Basidiomycota*) recorded from the Brazilian Atlantic Forest. *Mycotaxon* 109: 439–442. doi:10.5248/109.439
- Bononi VLR. 1979. Basidiomicetos do parque Estadual da Ilha do Cardoso: III. Espécies hidnóides. *Rickia* 8: 63–74.
- Bononi VLR. 1981. Alguns basidiomicetos hidnóides da região Amazônica. *Rickia* 9: 13–30.
- Bononi VLR, Trufem SFB, Grandi RAP. 1981. Fungos macroscópicos do parque Estadual das Fontes do Ipiranga, São Paulo, Brasil, depositados no herbário do Instituto de Botânica. *Rickia* 9: 37–53.
- Bononi VLR, Oliveira AKM, Quevedo JR, Gugliotta AM. 2008. Fungos macroscópicos do Pantanal do Rio Negro, Mato Grosso do Sul, Brasil. *Hoehnea* 34(4): 489–511.
- Capelari M, Maziero R. 1988. Fungos macroscópicos no estado de Rondônia, região dos rios Jaru e Ji Paraná. *Hoehnea* 15: 28–36.
- Dennis RWG. 1970. Fungus flora of Venezuela and adjacent countries. *Kew Bulletin Additional Series* 3: 1–530.
- Eriksson J, Ryvarden L. 1976. The *Corticaceae* of North Europe, vol. 4. *Hyphodermella–Mycoacia*. Fungiflora, Oslo, Norway.
- Gibertoni TB, Santos PJP, Cavalcanti MAQ. 2007. Ecological aspects of *Aphyllorales* in the Atlantic Rain Forest in Northeast Brazil. *Fungal Diversity* 25: 49–67.

- Gilbertson RL, Ryvarde L. 1986. North America Polypores, vol.1. *Abortiporus-Lindtneria*. Fungiflora, Oslo, Norway.
- Hjortstam K, Bononi VLR. 1986a. Fungos corticídeos do Brasil (*Aphyllphorales*). Rickia 13: 113–125.
- Hjortstam K, Bononi VLR. 1986b. Studies in tropical *Corticaceae* (*Basidiomycetes*) VI: A new species of *Steccherinum* from Brazil. Mycotaxon 25: 467–468.
- Hjortstam K, Bononi VLR. 1987. A contribution to the knowledge of *Corticaceae* s.l. (*Aphyllphorales*) in Brazil. Mycotaxon 28: 1–15.
- Hjortstam K, Ryvarde L. 1996. New and interesting wood-inhabiting fungi (*Basidiomycotina-Aphyllphorales*) from Ethiopia. Mycotaxon 60: 181–190.
- Ipulet P, Ryvarde L. 2005. New and interesting polypores from Uganda. Synopsis Fungorum 20: 87–99.
- Jarenkow JA, Budke JC. 2009. Padrões florísticos e análise estrutural de remanescentes florestais com *Araucaria angustifolia* no Brasil. 113–126, in: Fonseca CR, Souza AF, Leal-Zanchet AM, Dutra T, Backes A, Ganade G (Org.). Floresta com araucária: ecologia, conservação e desenvolvimento sustentável. Ribeirão Preto, Holos Editora.
- Jesus MA. 1993. Basidiomicetos lignocelulolíticos de floresta nativa e de *Pinus elliottii* Engelm. do Parque Estadual das Fontes do Ipiranga, São Paulo, SP. Hoehnea 20(1/2): 119–126.
- Nietiedt SA, Guerrero RT. 2000. Família *Corticaceae* (s.l.) – Novas citações para o Brasil. Iheringia sér. Bot. Porto Alegre 54: 45–56.
- Oliveira-Filho AT, Jarenkow JA, Budke JC. 2009. Patterns of floristic differentiation among Atlantic forests and the influence of climate II: The southern scenario. In: Impacts of Global Changes on Tropical Ecosystems Cross-cutting the Abiotic, Biotic and Human Spheres. Joint Meeting of the Association for Tropical Biology and Conservation (ATBC) and Society for Tropical Ecology, Marburg.: Philipps Universität Marburg, p. 433-433.
- Piatek M, Seta D, Szczepkowski A. 2004. Notes on Polish polypores 5. Synopsis of the genus *Spongipellis*. Acta Mycologica 39(1): 25–32.
- Rajchenberg M, Meijer AAR. 1990. New and noteworthy polypores from Paraná and São Paulo States, Brazil. Mycotaxon 38: 173–185.
- Reid DA. 1976. Notes on polypores. 2. Memoirs of the New York Botanical Garden 28(1): 179–198.
- Rick J. 1932a. Hidnaceae Riograndenses. Egatea 17: 1–9.
- Rick J. 1932b. Hidnaceae Riograndenses. Egatea 17(4–5): 209–212.
- Rick J. 1959. Basidiomycetes eubasidii in Rio Grande do Sul – Brasília. 3. *Hypochnaceae*, *Clavariaceae*, *Craterellaceae*, *Hydnaceae*. Iheringia, Bot. 5: 125–192.
- Ryvarde L. 1991. Genera of polypores. Nomenclature and taxonomy. Synopsis Fungorum 5: 1–363.
- Sonego RC, Backes A, Souza AF. 2007. Tree community structure and composition in a mixed ombrophilous forest in São Francisco de Paula, Rio Grande do Sul State, Brazil. Acta Botanica Brasílica 21(4): 943–955.
- SOS Mata Atlântica/INPE. 2009. [continuously updated]. Atlas dos Remanescentes da Mata Atlântica. <http://www.sosmatatlantica.org.br> [Accessed October 28, 2010].
- Sótão HMP, Bononi VLR, Figueiredo TS. 1991. Basidiomycetes de manguezais da Ilha de Maracá, Amapá, Brasil. Boletim do Museu paraense Emílio Goeldi, sér. Bot. 7(1): 109–114.
- Thiers B. 2007. [continuously updated]. Index Herbariorum. New York Botanical Garden, New York. <http://sciweb.nybg.org/science2/IndexHerbariorum.asp> [Accessed October 28, 2010].